

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search	PubMed	for				Go	Clear
		Limits	Preview/Index	History	Clipboard		
About Entrez		Display	Abstract	Save	Text	Order	Add to Clipboard

Entrez PubMed
Overview
Help | FAQ
Tutorial **NEW**
New/Noteworthy

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Cubby

Related Resources
Order Documents
Grateful Med
Consumer Health
Clinical Alerts
ClinicalTrials.gov

Privacy Policy

☐ 1: J Cell Biochem 1997 Feb;64(2):258-72

Related Articles, Books, LinkOut

InterScience

Purification and substrate specificity of polydeoxyribonucleotide kinases isolated from calf thymus and rat liver.


Karimi-Busheri F, Weinfeld M.

Experimental Oncology, Cross Cancer Institute, Edmonton, Alberta, Canada.

Damage to DNA can result in strand breaks with 5'-hydroxyl and 3'-phosphate termini. Before DNA polymerases and ligases can rejoin the broken strands, such termini have to be restored to 5'-phosphate and 3'-hydroxyl groups. Polydeoxynucleotide kinase is an enzyme that may fulfil this function. We have purified the kinases from calf thymus and rat liver to near homogeneity. Based on SDS-polyacrylamide gel electrophoresis and activity gels, the enzymes from both sources are approximately 60-kDa polypeptides. Both enzymes have an acidic pH optimum (5.5-6.0) for kinase activity, and similar pI values (8.5-8.6), and a specificity for DNA. The calf thymus kinase possesses a 3'-phosphatase activity, as has previously been shown for the rat liver enzyme. The minimum size of oligonucleotide that can be labelled is 7-8 nucleotides in length, but the optimal size appears to be > 18 nucleotides. Comparison of phosphorylation of oligo(dA)₂₄ and oligo(dT)₂₄ with oligonucleotides containing a varied nucleotide sequence indicated that the homopolymers are poorer substrates. Unlike the bacteriophage T4 polynucleotide kinase, the mammalian kinases exhibit no preference for 5'-overhanging termini when acting at DNA termini produced by restriction enzymes. With double-stranded oligonucleotide complexes designed to mode single-strand gaps and nicks, the mammalian kinases preferentially phosphorylate the 5'-terminus associated with the gap or nick, in keeping with the idea that the kinases are involved in the repair of DNA single-strand breaks.

PMID: 9027586 [PubMed - indexed for MEDLINE]

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	
Search	PubMed	for					Go	Clear
		Limits	Preview/Index	History	Clipboard			

[About Entrez](#)**Entrez PubMed**[Overview](#)[Help | FAQ](#)[Tutorial](#)[New/Noteworthy](#)**PubMed Services**[Journal Browser](#)[MeSH Browser](#)[Single Citation Matcher](#)[Batch Citation Matcher](#)[Clinical Queries](#)[Cubby](#)**Related Resources**[Order Documents](#)[Grateful Med](#)[Consumer Health](#)[Clinical Alerts](#)[ClinicalTrials.gov](#)[Privacy Policy](#)☐ 1: J Cell Biochem 1995 May;58(1):115-31[Related Articles, Books, LinkOut](#)**5'phosphorylation of DNA in mammalian cells: identification of a polymin P-precipitable polynucleotide kinase.****Prinos P, Slack C, Lasko DD.**

Department of Medicine, McGill University, Montreal, Quebec, Canada.

Proteins that catalyze 5' phosphorylation of an oligodeoxyribonucleotide substrate can be fractionated by polymin P treatment of whole cell extracts of calf thymus glands. Anion exchange chromatography on Q-Sepharose revealed three separable peaks of activity in the polymin P supernatant fraction, and one peak of activity in the Polymin P pellet fraction. The latter activity, Polymin P-precipitable polynucleotide kinase (PP-PNK), was further purified with a 1,500-fold increase of specific activity compared to the crude Polymin P pellet fraction. Oligonucleotides, a dephosphorylated 2.9-kb EcoRI fragment, and poly (A) were phosphorylated by the enzyme preparation, but thymidine 3' monophosphate was not a substrate. PP-PNK preparations exhibited an apparent K_M of 52 μM for ATP and 8 μM for oligo dT25. The enzyme preparation displayed no detectable 3' phosphatase or cyclic 2',3' phosphohydrolase activities. The sedimentation coefficient of the PP-PNK activity was 3.8S as determined by sucrose density gradient analysis; the Stokes radius was 45 Å, leading to an estimated molecular mass of 72 kDa. The enzyme had a pH optimum in the neutral to alkaline range in several buffer systems and is distinct from the DNA kinase with an acidic pH optimum previously described in calf thymus.

PMID: 7642718 [PubMed - indexed for MEDLINE]

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer